

A Multi-Phase Chemo-Dynamical SPH Code for Galaxy Evolution. Testing the Code.

Peter Berczik (berczik@mao.kiev.ua)

Main Astron. Obs. of Ukrainian National Academy of Sciences, Kiev, Ukraine

Gerhard Hensler and Christian Theis

Institut für Theoretische Physik und Astrophysik, University of Kiel, Germany

Rainer Spurzem

Astronomisches Rechen-Institut, Heidelberg, Germany

Abstract. In this paper we present some test results of our newly developed Multi-Phase Chemo-Dynamical Smoothed Particle Hydrodynamics (MP-CD-SPH) code for galaxy evolution. At first, we present a test of the “pure” hydro SPH part of the code. Then we describe and test the multi-phase description of the gaseous components of the interstellar matter. In this second part we also compare our condensation and evaporation description with the results of a previous 2d multi-phase hydrodynamic mesh code.

Keywords: multi-phase SPH, chemo-dynamical code, galaxy evolution

1. Introduction

Since several years Smoothed Particle Hydrodynamics (SPH) (Monaghan, 1992) has been applied successfully to study the formation and evolution of galaxies. Its Lagrangian nature as well as its easy implementation together with standard N-body codes allows for a simultaneous description of complex dark matter-gas-stellar compositions (Navarro & White, 1993; Mihos & Hernquist, 1996). Nevertheless, until now the codes lack processes that are based on the coexistence of different phases of the interstellar medium (ISM), mainly dissipative, dynamical and stellar feedback, element distributions, etc. We have therefore developed a 3d chemo-dynamical code which is based on our single phase galactic evolutionary program (Berczik, 1999; Berczik, 2000). This code includes many complex effects such as a multi-phase ISM, cloud-cloud collisions, the drag force between different ISM components, condensation and evaporation of clouds (CE), star formation (SF) and stellar feedback (FB). The more detailed description of the new code will be presented in a comprehensive paper (Berczik et al., in preparation).



© 2008 Kluwer Academic Publishers. Printed in the Netherlands.

2. The hydro code test

The self-gravitating collapse of an initially isothermal, cool gas sphere is a common test problem for SPH codes (Evrard, 1988; Hernquist & Katz, 1989; Steinmetz & Müller, 1993; Carraro et al., 1998; Thacker et al., 2000; Springel et al., 2001). Following these authors, we consider a gas sphere of total mass M , radius R , and initial density profile $\rho(r) = \frac{M}{2\pi R^2} \frac{1}{r}$ and with an internal energy per unit mass of $u = 0.05 \frac{GM}{R}$. At the beginning, the gas particles are at rest. We use a system of units with $G=M=R=1$. In Fig. 1. the time evolution of the different energies (left) and of the relative error of the total energy (right) are displayed. In the right figure we also compare results applying the publicly accessible **GADGET** code (Springel et al., 2001) with “standard” parameters for the 32,000 particles. During the central bounce around $t \approx 1.1$ most of the kinetic energy is converted into heat and a strong shock wave travels outwards. The results of Fig. 1. agree very well with those of Steinmetz & Müller (1993) and Springel et al. (2001). The maximum relative total energy error is around 0.05 % even for low (8,000) particle numbers.




Figure 1. Time evolution of the thermal, kinetic, potential and total energy for the collapse of an initially isothermal gas sphere (left) and the relative total energy error for different models (right). The different lines correspond to the different numbers of gas particles. In the right figure we also present the energy error of the **GADGET** code with “standard” parameters for the 32,000 particles.

3. The multi-phase processes

In our multi-phase gas code we use a two-component gas description of the ISM (Theis et al., 1992; Samland et al., 1997), a cold (10^2 - 10^4 K)

cloudy component treated by sticky particles (Theis & Hensler, 1993) and a diffuse hot gas (10^4 - 10^7 K) represented by SPH particles. The cold clumps follow an empirical mass-radius relation (Larson, 1981; Solomon et al., 1987; Maloney, 1990; Inoue & Kamaya, 2000; Theis & Hensler, 1993) with $h_{cl} \simeq 50\sqrt{m_{cl}/(10^6 M_\odot)}$ pc and experience dissipation due to cloud-cloud collisions and drag forces within the hot gas. The mass exchange between “cold” and “hot” gas phases happens basically through condensation and/or evaporation (CE) described by Cowie et al. (1981) and Köppen et al. (1998). The basic parameter switching between both processes, σ_0 , is set to 0.03 according to Cowie et al. (1981).




Figure 2. The time evolution of the mass exchange rate due to condensation *vs.* evaporation (CE) effects (left). The letters indicate the phase (i.e. H - Hot; C - Cold) and the numbers indicate the different models. In the right figure we compare our “best” model with 2d multi phase hydro mesh code results, for similar initial conditions.

As a CE test for our code, we calculate the evolution of a “hypothetical” gas system inside a dark matter (DM) halo and compare the results with a 2d multi-phase hydro mesh-code (Rieschick & Hensler, 2000) for the same system. The initial total gas mass amounts to $2 \times 10^9 M_\odot$ (99% “COLD” + 1% “HOT”) and is distributed according to a Plummer-Kuzmin disk with parameters $a=0.1$ kpc and $b=2$ kpc (Miyamoto & Nagai, 1975). The initial temperature for the cold gas is set to 2×10^3 K, for the hot gas to 10^6 K. The DM halo is static with parameters $r_0=2$ kpc and $\rho_0 = 2 M_\odot/\text{pc}^3$ (Burkert, 1995) so that its mass inside the initial gas distribution (20 kpc) amounts to $\simeq 3.2 \times 10^{11} M_\odot$.

In some of the models we also use an initial cloud mass function (ICMF) (Miyazaki & Tsuboi, 2000) with a power-law index -1.5 and lower and upper mass limits at $10^4 M_\odot$ and $10^6 M_\odot$, respectively. In

Fig. 2. we present the time evolution of the mass exchange rate via CE for different models: **VER-01** with 5,000 hot-gas SPH particles of identical mass and 5,000 cold gas particles, **VER-02** with 10,000 particles each, **VER-03** with the same particle numbers as in **VER-01** but an ICMF, and **VER-04** equal to **VER-02** but also with an ICMF. The figures show that the basic behaviour of mass exchange between the hot and cold phase is well described with our model and almost independent of the particle number and ICMF, but in agreement on average with the rates of the 2d mesh code.

Acknowledgements

The work was supported by the German Science Foundation (DFG) under grant *436 UKR* and by the *SFB 439* at the University of Heidelberg. P.B. is grateful for the hospitality of the Inst. für Theor. Physik und Astrophysik in Kiel where most of this work was done. The calculations have been performed with the **GRAPE5** system at the Astronomical Data Analysis Center of the NAO, Japan.

References

- Berczik P., 1999, *A&A*, **348**, 371
 Berczik P., 2000, *Ap&SS*, **271**, 103
 Burkert A., 1995, *ApJ*, **447**, L25
 Carraro G., Lia C. & Chiosi C., 1998, *MNRAS*, **297**, 1021
 Cowie L.L., McKee C.F. & Ostriker J.P., 1981, *ApJ*, **247**, 908
 Evrard A. E., 1988, *MNRAS*, **235**, 911
 Hernquist L. & Katz N., 1989, *ApJ*, **70**, 419
 Inoue A.K. & Kamaya H., 2000, *PASJ*, **52**, L47
 Köppen J., Theis Ch. & Hensler G., 1998, *A&A*, **331**, 524
 Larson R.B., 1981, *MNRAS*, **194**, 809
 Maloney P., 1990, *ApJ*, **349**, L9
 Mihos J.C. & Hernquist L., 1996, *ApJ*, **464**, 641
 Miyamoto M. & Nagai R., 1975, *PASJ*, **27**, 533
 Miyazaki A. & Tsuboi M., 2000, *ApJ*, **536**, 357
 Monaghan J.J., 1992, *ARA&A*, **30**, 543
 Navarro J.F. & White S.D.M., 1993, *MNRAS*, **265**, 271
 Rieschick A. & Hensler G., 2000, *ASP Conf. Ser.*, **215**, 130
 Samland M., Hensler G. & Theis Ch., 1997, *ApJ*, **476**, 544
 Solomon P.M., Rivolo A.R., Barrett J. & Yahil A., 1987, *ApJ*, **319**, 730
 Springel V., Yoshida N. & White S.D.M., 2001, *NewA*, **6**, 79
 Steinmetz M. & Müller E., 1993, *A&A*, **268**, 39
 Thacker R. J., Tittley E. R., Pearce F. R., et al., 2000, *MNRAS*, **319**, 619
 Theis Ch., Burkert A. & Hensler G., 1992, *A&A*, **265**, 465
 Theis Ch. & Hensler G., 1993, *A&A*, **280**, 85